

# Temperature Effects on Greenhouse Gas Production From Treatment Wetland Soils Along a Nutrient Gradient

By: Kyle Dittmer and Havalend Steinmuller

Faculty Mentor: Dr. Lisa Chambers

UCF Department of Biology

**ABSTRACT:** It is generally accepted that increased temperatures are positively correlated with microbial respiration rates, causing greater greenhouse gas (GHG) emissions ( $\text{CO}_2$  and  $\text{CH}_4$ ) from wetlands. The goal of this study was to understand the interacting effects of temperature and nutrient concentrations on GHG emissions from wetland soils. Complementary field studies and a laboratory study were completed within Cell 1 of the Orlando Wetlands Park (Christmas, FL). Four sampling locations were established along a transect and sampled in summer and winter for the field studies. Soils (0-10 cm) were incubated under anaerobic conditions for 48-hours at ambient or elevated temperatures and GHG flux was measured. Surface water nutrients and soil physiochemical properties were also analyzed. Carbon dioxide and methane production both differed through an interaction between season and site ( $p = 0.04$  and  $p < 0.001$ , respectively) with higher rates in the summer and at sites with the higher soil organic matter; water nutrients did not have a significant effect on GHG emissions. The laboratory study used only one soil sample, but varied water nutrients and temperature (3 x 3 factorial design) using four replicates per treatment and incubating under anaerobic conditions for 10 days. Temperature had a significant effect on both  $\text{CO}_2$  and  $\text{CH}_4$  production (both  $p < 0.001$ ), but water nutrients did not have an effect, presumably due to exiting high nutrient levels within the soil porewater. These results highlight the importance of soil properties (organic matter content and porewater nutrients) when determining the influence of temperature and water nutrients on GHG production.

**KEYWORDS:** wetland; treatment wetland; greenhouse gasses; eutrophication; climate change; biogeochemistry; soil

## 1. INTRODUCTION

Excess amounts of greenhouse gasses (GHGs), particularly carbon dioxide (CO<sub>2</sub>) and methane (CH<sub>4</sub>), alter the energy balance of the Earth's climate system and are major drivers of climate change. Since the beginning of the Industrial Revolution, atmospheric CO<sub>2</sub> has increased from 278 ppm to 406 ppm (NOAA, 2017), while CH<sub>4</sub> has increased from 722 ppb to 1803 ppb throughout the same time scale (Ciais et al., 2013; Ballantyne et al., 2012). The largest proportion of these emissions are from anthropogenic manipulations, such as the burning of fossil fuels and land use change (IPCC, 2007; Johnson et al., 2007). Carbon dioxide accounts for approximately 65% of GHGs in the atmosphere. Although CH<sub>4</sub> accounts for a smaller percentage, it absorbs 30 times the infrared radiation absorbed by CO<sub>2</sub>, contributing to increased warming (Houghton, 2001).

Greenhouse gas emissions from wetlands are primarily mediated by microbial respiration and decomposition. Wetlands are often inundated or saturated, causing microbially-mediated decomposition rates to be comparatively slower than adjacent upland environments (Reddy and DeLaune, 2008). This lack of oxygen, the most preferred electron acceptor for microorganisms, causes facultative aerobic bacteria and anaerobic bacteria to utilize alternative electron acceptors, resulting in different end products (Table 1). The rate at which these alternate electron acceptors are consumed depends on their relative abundance, the quantity of organic compounds (electron donor), and the microbial populations present (Reddy and DeLaune, 2008).

Many scientists expect increasing temperatures resulting from climate change will increase soil microbial respiration rates (Brooks et al., 1997; Butterbach-Bahl et al., 2011; Dalal and Allen, 2008; Holst et al., 2008). This increase in respiration rates is particularly important in wetland soils, which serve as a vast reservoir of accumulated carbon (C), nitrogen (N), and phosphorus (P) by incorporating dead plant biomass and other detritus into the soil matrix (Larmola et al., 2013). If warmer temperatures accelerate decomposition in wetland soils, a greater efflux of CO<sub>2</sub> and CH<sub>4</sub> is expected, creating a positive GHG feedback loop that could promote further warming.

Concurrently, anthropogenic activities, predominantly agricultural fertilizer use, cause eutrophication (bodies of water receiving excess nutrients). Most notable are N and

P, primarily due to human activities such as agricultural fertilizer use and point-source pollution (Webber, 2010). This excess of typically limiting nutrients promotes the prolific growth of algae and other vegetation, the potential for harmful algal blooms, a decrease in water transparency, changes in fish communities, and hypoxic waters that result in "dead zones" (Ansari, 2011). As the human population continues to rise (increasing from 6.5 billion people in 2005 to a projected 9.2 billion in 2050 (UNPD, 2008)), fertilizer use is predicted to increase by 40 percent between 2002 and 2030 (FOA, 2000), thus contributing to amplified eutrophication. Nitrate (NO<sub>3</sub><sup>-</sup>), ammonium (NH<sub>4</sub><sup>+</sup>), and soluble reactive phosphorous (SRP) are the most bioavailable forms of N and P and greatly contribute to anthropogenic eutrophication (McCormick and Laing, 2003).

Treatment wetlands are constructed ecosystems dominated by aquatic vegetation that use natural processes to remove pollutants (Reddy and DeLaune, 2008). These systems can mitigate anthropogenic eutrophication by utilizing wetland vegetation, soils, and associated microbial communities to improve water quality (Chen, 2001; Kadlec, 2008, 2009; Marimon and Chang, 2016). They are commonly constructed for several different purposes, such as treating anthropogenic wastewater (domestic, agricultural, and industrial), reducing metals, and treating pathogens by biological means (Kadlec, 2009) at an inexpensive cost. However, it is not fully understood how both temperature and bioavailable nutrient concentrations interact to affect GHG emissions from treatment wetland soils. The goal of this study is to scrutinize the effect of climate change scenarios (increased temperatures and eutrophication) on CO<sub>2</sub> and CH<sub>4</sub> production in treatment wetland soils in order to assess the role of treatment wetlands to future GHG emissions. We hypothesized that GHG flux will be positively correlated with increased temperature and bioavailable nutrient concentrations.

## 2. METHODS

### 2.1 Site Description

The Orlando Wetlands Park (OWP), located in Christmas, Florida, spans 1,220 acres (Kadlec and Wallace, 2008). The OWP is designed to use wetland macrophytes to promote N and P removal and polish off tertiary treated domestic wastewater from the City of Orlando before entering the St. Johns River (Sees and Turner 1997). The Florida Department of Environmental

Protection (FDEP) permits annual discharge limits of 2.31 mg L<sup>-1</sup> for total nitrogen (TN) and 0.200 mg L<sup>-1</sup> for total phosphorous (TP). In 2015, the OWP's annual discharge concentration for TN and TP were 0.95 mg L<sup>-1</sup> and 0.075 mg L<sup>-1</sup>, respectively, demonstrating the functionality of the constructed wetland (Rothfeld, 2015).

The OWP is divided into seventeen cells and three distinct wetland communities: deep marsh, mixed marsh, and a 90-acre lake. The deep marsh encompasses cells 1–12 and is strategically planted with cattails (*Typha latifolia*) and bulrush (*Scirpus spp.*) due to their ability to take up nutrients in wastewater (US. EPA. 1993; Figure 1). Average water depth of the deep marsh is 1 to 1.5 meters and is inundated year-round. The OWP is an excellent location to test our hypotheses due to the clear nutrient gradient that develops as water flows throughout the park from the inflow to the outflow. Based on data from the park's management, Stratum 1 (Cells 1, 2, 11, and 12) has been shown to be the most efficient system at polishing off NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup>, and SRP within the surface water. Cell 1 served as our sampling location, as it contained an influent structure (source of nutrient-rich water entering the OWP), which served as a proxy for a eutrophic system, and because it was a part of the most successful stratum, meaning a large nutrient gradient should be present.

## 2.2 OBSERVATIONAL FIELD STUDY

### 2.2.1 Soil Sampling

We established four sampling sites along a 660 meter transect spanning the length of Cell 1 (Figure 1), spaced ~165 m apart. Sampling locations contained similar vegetation, including cattail (*Typha latifolia*) and bulrush (*Scirpus spp.*). Three soil cores were taken at each of the four sites via the push-core method, and each soil core was extruded in the field into two depth intervals: 0–10 cm and 10–20 cm. Soil samples were placed in sealed polyethylene bags. Surface water samples (1 L) were collected at each site in acid-washed polyethylene bottles to determine ambient concentrations of NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, and SRP via U.S. EPA methods 50.1 Revision 2.0, 353.2 Revision 2.0, and 365.1 Revision 2.0, respectively (U.S. EPA, 1993). Sampling occurred twice along this transect (June 27, 2016 and January 20, 2017) to capture seasonal variability. Segmented samples were placed on ice and immediately transported back to the laboratory, where field replicates were homogenized to provide a

representative sample of site conditions at each depth.

### 2.2.2 Greenhouse Gas Production (CO<sub>2</sub> and CH<sub>4</sub>)

Eight (8) replicate bottle incubations were created within 24 hours of sample collection by weighing 15 g field-moist soil, exclusively from the 0–10 cm depth interval, into 100 mL serum bottles. The bottles were capped with a rubber septa and aluminum crimp, evacuated to -75 mmHg, and purged with 99% O<sub>2</sub>-free N<sub>2</sub> gas for three minutes in order to mimic the anaerobic conditions of the site. Fifteen (15) mL of site water was filtered via Supor 0.45 μm membrane filters, purged with N<sub>2</sub> gas, and injected into each bottle. Four bottles from each site were randomly assigned to each of two treatments—ambient and elevated temperature. Ambient temperature was defined as the atmospheric temperature during the time of sampling; elevated temperature was based on a four degree Celsius increase from the ambient temperature. The four degree Celsius increase was chosen for the elevated temperature due to the IPCC report "Climate Change 2007: Impacts, Adaptation, and Vulnerability," which concludes that the global average temperature is predicted to increase 1.4–5.8°C by 2100 (Parry et al., 2007). During the summer sampling, ambient temperature was 32°C and elevated was 36°C. During winter, ambient temperature was 22°C and elevated was 26°C. Bottles were incubated continuously in an orbital shaker at 150 rpm at their respective temperature treatment for 48 hours. Carbon dioxide and CH<sub>4</sub> samples were extracted from the bottle headspace using gas-tight syringes at 6, 12, 24, and 48 hours and analyzed using a GC-2014 Gas Chromatograph (Shimadzu Instruments, Kyoto, Japan) equipped with a flame ionization detector (FID), and a Hayesep N 80/100 Mesh 1/8in X 1.5M stainless pre-conditioned column. Breathing grade air was used as the carrier gas. The temperature of the injection port was sufficient to sterilize the syringe between injections. Respiration rates at each temperature were calculated as linear slopes obtained by repeated measures of CO<sub>2</sub> and CH<sub>4</sub> concentrations over time. The fraction of CO<sub>2</sub> and CH<sub>4</sub> dissolved in the liquid phase was calculated via the use of Henry's Law.

## 2.3 CONTROLLED LABORATORY STUDY

### 2.3.1 Soil Sampling

A laboratory study was conducted to determine the interactive effects of temperature and nutrient concentrations on greenhouse gas emissions, independent

of other environmental variables. Fourteen (14) cores were collected within a 5 m<sup>2</sup> area in Cell 1 (28°34'20.2"N and 81°00'33.2"W) and extruded to a depth of five (5) centimeters. Soils were stored in sealed polyethylene bags on ice and immediately transported to the laboratory, where samples were homogenized to eliminate spatial variability. In addition to soil cores, 1 L of surface water was collected in an acid-washed polyethylene bottle.

### 2.3.2 Greenhouse Gas Production (CO<sub>2</sub> and CH<sub>4</sub>)

To determine the interactive effects of nutrient concentrations and temperature on greenhouse gas emissions, soil bottles were randomly assigned to one of three temperature treatments and also one of three nutrient treatments in a 3 x 3 factorial design. Again, there were four replicates per treatment, totaling 36 bottles. The same preparations for bottle incubations were followed for the controlled laboratory study as described in section 2.2.2 of this paper. Three temperature treatments were established (25°C, 30°C, and 35°C) and were randomly assigned to one of three different aqueous nutrient treatments: high, medium, or low concentrations of ammonium, phosphate, and glucose. Ammonium was chosen as the N source because it is the predominate form available in anaerobic wetland soils. Phosphate was used for the P source because it is the bioavailable form of P (Reddy and DeLaune, 2008). To isolate the effects of the N and P treatments and avoid impeding microbial metabolism due to C limitations, all three nutrient treatments received a non-limiting amount of glucose as a C source. The phosphate solution was created using 1000 ppm Phosphate Standard and the ammonium solution was made from Ammonia Standard, 1000 ppm NH<sub>3</sub>, both by Ricca Chemical Company. The glucose solution was made from D(+)-Glucose monohydrate (manufactured by Acros Organics) from Fisher Scientific (Table 2). The 'High' nutrient concentrations were defined as the highest level of the respective nutrient that has ever been loaded into the OWP, which was 2.2 mg L<sup>-1</sup> and 0.5 mg L<sup>-1</sup> for TN and TP, respectively (Rothfeld, 2015). 'Medium' nutrient concentrations were half of the high concentrations; 'low' nutrient concentrations received pure nanopure water (Table 2). All four replicates per treatment were incubated continuously in an orbital shaker at 150 rpm at their respective temperatures for ten days, while CO<sub>2</sub> and CH<sub>4</sub> production was measured at intervals of 24, 48, 72, 144, 192, and 240 hours.

### 2.4 Soil and Water Properties

Within the laboratory, the chemical and physical properties of the soils at each depth segment from both sections 2.2.1 and 2.3.1 were analyzed, including bulk density, percent organic matter (OM), percent moisture content, pH, extractable nutrients, total carbon (TC), TN, and TP. Surface water and porewater were filtered through Supor 0.45 µM filters and acidified with double-distilled sulfuric acid (DDI H<sub>2</sub>SO<sub>4</sub>) for preservation and analyzed for NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup>, and SRP concentrations. Samples were stored at 4°C and all analyses were performed within three weeks of sampling.

Moisture content and bulk density were determined by drying a subsample at 70°C for three days until a constant weight was achieved. Dried soils were ground using a SPEX Sample Prep 8000M Mixer/Mill (Metuchen, NJ). Ground soils were analyzed for TC and TN using a Vario Micro Cube CHNS Analyzer (Elementar Americas Inc., Mount Laurel, NJ). Percent OM was determined by loss on ignition (LOI), while dry soils were combusted at 550 °C for 5 hours and final weight was subtracted from initial weight. Total P was determined by digesting the resulting ash from LOI in 50 mL of 1 N HCl on a hot plate for 30 minutes at 65.5 °C. The TP solution was then filtered through a Whatman #41 filter paper (Andersen, 1976). Extractable porewater nutrients (NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, and SRP) were processed by placing 2.5 g of wet soil into a 40 mL centrifuge tube with 25 mL of 1 M KCl and shaking the solution for an hour at 150 rpm. Samples were then placed in a centrifuge for 10 minutes at 4000 rpm and 10°C, filtered through Supor 0.45 µM filters, and acidified with DD H<sub>2</sub>SO<sub>4</sub> for preservation. Surface water, porewater, TP, and extractable nutrients samples were analyzed using an AQ2 Automated Discrete Analyzer (Seal Analytical, Mequon, WI) according to EPA method 365.1 Rev. 2 (U.S. EPA, 1993).

### 2.5 Microbial Biomass Carbon

Microbial Biomass C (MBC) was determined in all depth segments by chloroform fumigation (Vance et al., 1987). Fumigate and non-fumigate duplicates were prepared by placing 2.5 g of wet soil in 40 mL centrifuge tubes. The fumigated samples were exposed to volatilized chloroform for 24 hours in a glass vacuum desiccator in order to detect the organic C within the microbial cell wall. After 24 hours, the fumigated samples were extracted with 25 mL of 0.5M K<sub>2</sub>SO<sub>4</sub>, placed in an orbital shaker for 1 hour at 150 rpm and 25°C, then

centrifuged for 10 minutes at 4000 rpm and 10°C. The sample was vacuum-filtered through Supor 0.45  $\mu\text{m}$  filters, and acidified with DD  $\text{H}_2\text{SO}_4$  for preservation. The non-fumigate samples served to quantify dissolved organic C and were prepared in the same way, except for not being exposed to the chloroform fumigation. Total organic carbon (TOC) was found by using a Shimadzu TOC-L Analyzer (Kyoto, Japan) for both treatments. Microbial biomass C was calculated by the difference between the fumigated samples and the non-fumigated samples.

### 2.6 Statistics

In order to determine the temperature and nutrient effects on GHG production, statistical analysis was conducted in R (R Foundation for Statistical Computing, Vienna, Austria). Datasets met the assumptions of homogeneity of variance, which was tested via use of the Levene's test ( $\alpha = 0.05$ ). Normality was verified by the Shapiro-Wilks test ( $\alpha = 0.05$ ); datasets that did not meet the assumptions of normality were logarithm-transformed to verify normality. A two-way ANOVA was used to determine the significance of predictor variables temperature, season, and site (distance from inflow) in predicting  $\text{CO}_2$  and  $\text{CH}_4$  production ( $\alpha = 0.05$ ). Akaike's Information Criterion (AIC) was used to implement a model simplification approach, which determined that the most appropriate predictor variables were season, site (distance) and the interaction between them for the field study. The same approach was implemented in the controlled laboratory study, which determined that temperature was the only significant predictor variable to determine GHG emissions, which excluded nutrient availability. Tukey post-hoc tests were used to determine significance of pairwise comparisons, also using  $\alpha = 0.05$ .

## 3. RESULTS

### 3.1.1 Observational Field Study Greenhouse Gas Production

Carbon dioxide production differed by season and site ( $p < 0.001$ ,  $p = 0.052$ ), as well as the interaction between the two predictor variables ( $p = 0.043$ ).  $\text{CO}_2$  production ranged from  $0.226 \pm 0.059 \mu\text{g CO}_2\text{-C g}^{-1} \text{d}^{-1}$  at site 4 during the winter to  $17.3 \pm 7.23 \mu\text{g CO}_2\text{-C g}^{-1} \text{d}^{-1}$  at site 2 during the summer (Figure 2). In general, summer  $\text{CO}_2$  production was greater than winter  $\text{CO}_2$  production. Though site was a significant predictor of  $\text{CO}_2$  production, a clear trend of decreasing  $\text{CO}_2$  production with increasing distance from the inflow was

only evident during the winter sampling. Temperature did not have a significant effect on  $\text{CO}_2$  production, and was thus not incorporated into the model, but  $\text{CO}_2$  production was positively correlated to  $\text{CH}_4$  production, TN, TC, MBC, percent OM, and moisture content, while  $\text{CO}_2$  production was negatively correlated to bulk density and C/N ratio (Table 3).

Similar to the results of  $\text{CO}_2$  production,  $\text{CH}_4$  production differed by both season and site, and there was a significant interaction between the two variables (all  $p < 0.001$ ). Sites 2 and 3 were not significantly different from each other; all other sites differed from each other. Winter  $\text{CH}_4$  production was lower than summer  $\text{CH}_4$  production. Methane production ranged from  $0.002 \pm 0.0001 \mu\text{g CH}_4\text{-C g}^{-1} \text{d}^{-1}$  during the winter at site 3 to  $0.411 \pm 0.030 \mu\text{g CH}_4\text{-C g}^{-1} \text{d}^{-1}$  at site 3 during the summer (Figure 3). Temperature treatment (ambient and elevated) was excluded from the model, as they were not significant predictor variables for  $\text{CH}_4$ . Methane was significantly correlated to  $\text{CO}_2$  production, TN, TC, MBC, percent OM, and moisture content, while negatively correlated to bulk density and C/N ratio (Table 3). In addition to soil physiochemical properties,  $\text{CH}_4$  production was significantly correlated to surface water  $\text{NH}_4^+$  and porewater  $\text{NH}_4^+$ .

Though there were observable differences in soil physiochemical properties and microbial biomass C (MBC) at each site, low replication prevented analysis via ANOVA due to the soils being homogenized for a representative sample. However, all soil physiochemical properties were correlated to each other (Table 3), as well as MBC. All correlations were positive except for bulk density, which was significantly negatively correlated to all other physiochemical properties and microbial biomass C.

### 3.1.2 Observational Field Study Nutrient Analysis

Surface water  $\text{NO}_3^-$  was highest at site 1 in winter with 1.13 mg N/L and lowest at sites 2 and 4, with no  $\text{NO}_3^-$  detected. Porewater  $\text{NO}_3^-$  was only detected in site 3 at 0.015 mg N/L in the winter and varied from 0.264 mg N/L at site 1 to 0.032 mg N/L at site 4 in the summer. Surface water  $\text{NO}_3^-$  varied from 0.136 mg N/L in site 1 to 0.011 mg N/L in site 3 during summer.

Surface water  $\text{NH}_4^+$  was found to be highest in site 2 with 0.163 mg N/L and lowest in site 1 with 0.091 mg N/L during the winter. Surface water  $\text{NH}_4^+$  increased

during the summer and varied from 0.338mg N/L in site 1 to 0.124 mg N/L at site 4. Porewater  $\text{NH}_4^+$  was highest in site 1 with 2.23 mg N/L and lowest in site 4 with 0.393 mg N/L in winter. During summer, there was a large spike in porewater  $\text{NH}_4^+$  which ranged from 12.4 mg N/L in site 1 to 3.07 mg N/L in site 4.

Surface water SRP showed an opposite trend during winter and had the highest concentration in site 4 (0.424 mg P/L) and the lowest in site 1 (0.270 mg P/L). In summer, surface water SRP was greatest within site 3 at 0.392 mg P/L and lowest in site 4 at 0.116 mg P/L. Winter porewater SRP was greatest in site 1 with 1.77 mg P/L and lowest in site 4 with 0.574 mg P/L. Summer porewater SRP was greatest in site 4 with 1.314 mg P/L and lowest in site 2 with 0.429 mg P/L.

Overall, the clear nutrient gradient we expected to find between the inflow and outflow of cell 1 was only observed for winter porewater SRP with nutrient concentrations decreasing with distance from the inflow. Other nutrients that had the highest concentrations in site 1 and lowest in site 4, but no linear decrease, were: winter surface water  $\text{NO}_3^-$ , winter surface water  $\text{NH}_4^+$ , summer surface water SRP, summer porewater  $\text{NO}_3^-$ , and summer and winter porewater  $\text{NH}_4^+$ . From this information, soil nutrients were determined to have a greater effect on GHG emissions from wetland soils compared to surface water nutrients because microbial populations cannot utilize the nutrients suspended within the surface water as readily.

### 3.2 Controlled Laboratory Study

Temperature treatments had a significant relationship with  $\text{CO}_2$  production ( $p < 0.001$ ). Carbon dioxide production increased with increasing temperature, ranging from  $0.966 \pm 0.037 \mu\text{g CO}_2\text{-C g}^{-1} \text{d}^{-1}$  at the lowest temperature treatment ( $25^\circ\text{C}$ ) to  $1.82 \pm 0.040 \mu\text{g CO}_2\text{-C g}^{-1} \text{d}^{-1}$  at the highest temperature treatment ( $35^\circ\text{C}$ ) (Figure 4a). Each temperature treatment differed from every other treatment. Temperature was also the only predictor of  $\text{CH}_4$  production ( $p < 0.001$ ), increasing with increasing temperature. The  $25^\circ\text{C}$  treatment produced  $0.149 \pm 0.011 \mu\text{g CH}_4\text{-C g}^{-1} \text{d}^{-1}$ , while the  $35^\circ\text{C}$  produced  $0.423 \pm 0.014 \mu\text{g CH}_4\text{-C g}^{-1} \text{d}^{-1}$  (Figure 4b). None of the concentrations of nutrients added had a significant effect on either  $\text{CO}_2$  or  $\text{CH}_4$  production, and thus were excluded from the model.

## 4. DISCUSSION

### 4.1 $\text{CO}_2$ and $\text{CH}_4$ Emissions

Several controlled laboratory studies and field studies have been conducted to determine the effects of temperature on GHG emissions (Brooks et al., 1997; Butterbach-Bahl et al., 2011; Chin et al., 1999; Dalal and Allen, 2008; Fang and Moncrieff, 2001; Holst et al., 2008; Inglett et al., 2012), while other studies have studied bioavailable nutrients as a factor for stimulating GHG emissions from soils (Bridgham and Richardson, 1992; Niu et al., 2010; Peng et al., 2011; Shi et al., 2014). The goal of our study was to use natural field conditions concomitant with a controlled laboratory study to determine if temperature and nutrient availability work synergistically to control GHG emissions from wetland soils. In general, temperature was positively correlated to  $\text{CO}_2$  and  $\text{CH}_4$  emissions, though the interaction with nutrient availability was less clear due to our experimental considerations (see Bioavailable Nutrients discussion below).

In a controlled laboratory study, Inglett et al. (2012) reported an increase in both emissions with increased temperature under aerobic and anaerobic conditions, using different soil types incubated at 10, 20, and  $30^\circ\text{C}$ . In our study, though not statistically significant, three out of the four sites had higher  $\text{CO}_2$  production at  $32^\circ\text{C}$  than at  $36^\circ\text{C}$ , suggesting that a maximum temperature threshold may occur that impedes microbial respiration. O'Connel (1990) also found in a laboratory study that the optimal temperature for microbial decomposition was approximately  $30^\circ\text{C}$  and any temperature above that threshold decreased respiration rates. However, this trend was not applicable to  $\text{CH}_4$  production, which suggests that methanogens (microbes that produce  $\text{CH}_4$  as a metabolic byproduct in anaerobic conditions) are more stimulated by increased temperatures compared to other soil microbes. After a 30-day incubation period of anoxic rice field soils, Chin et al. (1999) found that at  $30^\circ\text{C}$  methanogenic archaea predominated, whereas at  $15^\circ\text{C}$ , the archaeal community was significantly more diverse. These findings suggest that methanogenic microbes become more dominant as temperature continues to rise.

In a laboratory setting, Fang and Moncrieff (2001) found an exponential increase in  $\text{CO}_2$  emissions with increased temperature, up to  $32^\circ\text{C}$ , for upland soils with different soil moisture contents and incubation periods. This observation correlates to the findings of our laboratory study, suggesting that temperature is a

factor determining GHG flux rates, but there are more variables that regulate CO<sub>2</sub> production in wetland soils. Chapuis-Lardy et al. (2007) concluded that microbial activity, root respiration, soil organisms, and fungi were all sources of GHGs in soils. Additionally, Ludwig et al. (2001) found that soil moisture, pH, soil temperature, and bioavailable nutrients were also factors determining GHG emissions from soils.

Based on the significant correlations with OM and both GHG emissions in this study, our results suggest that one of the most important factors when considering GHG emissions from treatment wetland soils is the content of OM in the soils (Table 3). As OM content of the soils increases, microbial and plant communities will also increase in biomass due to the high bioavailable forms of C and N. In turn, as microbial communities increase in both biomass and activity, more soil OM will be decomposed and CO<sub>2</sub> and CH<sub>4</sub> will be produced at faster rates. By way of example, the soil from site 4 during the summer consisted of only 3.67% OM and thus had the lowest emissions for both GHGs under both ambient and elevated temperatures, while the highest soil OM content was found in site 3 (24.3%) and had the highest CH<sub>4</sub> emission and the second highest CO<sub>2</sub> emission for each temperature treatment. For the same season, site 4 also had the lowest microbial biomass (14,467.17 mg/kg) and site 3 had the highest (81,613.03 mg/kg). This trend was also found during the winter with site 1 having the highest soil OM content of 8.42% and the highest emissions for both GHGs under both temperature treatments. The lowest soil OM content for winter was in site 2 (1.58%) but had the second highest emissions for both GHG under both temperature treatments, although almost negligibly higher than sites 3 and 4 (Figure 2 and 3), which could potentially be explained by the sites' high moisture content (41.8%) (McKenzie et al., 1998).

The observed differences in soil OM along the field transect may be because Cell 1 was 'demucked' in 2001, where approximately 46 cm of sediment was removed in order to rejuvenate the hydraulic flow of surface water. We found that approximately 24 cubic yards of white builders sand was deposited and spread within Cell 1 in order to serve as an identifiable layer within the soil profile. The sand serves to quantify the rate of OM accumulation over time after the demucking procedure (personal communication, M. Sees). We found this sand layer approximately 18 to 25 cm below the surface layer for a majority of the samples. Additionally, the OWP has historically always had surficial soils consisting of fine

sand underlain by clayey soils (US. EPA. 1993). We suspect that this presence of fine sand is the primary reason why many of our soil samples contained minimal OM, compared to other freshwater marshes. Our results can be applicable to wetlands with low rates of vegetative senescence and thus slow soil OM accumulation rates, or to wetlands dominated by sandy soils.

#### 4.2 Bioavailable Nutrients

For the controlled laboratory study, we observed a general increase in CH<sub>4</sub> as nutrient concentrations and temperatures increased. Within the observational field study, porewater and surface water NH<sub>4</sub><sup>+</sup> were the only observed nutrients to be positively correlated with CH<sub>4</sub>. There is much discrepancy as to how NH<sub>4</sub><sup>+</sup> affects CH<sub>4</sub> emissions (Oertel et al., 2016). The negative correlation between the two can be explained at the biochemical level (Dunfield and Knowles, 1995; Gulledege and Schimel, 1998) and also at the microbial community level (Bodelier and Laanbroek, 2004). However, at the ecosystem level, increased availability of NH<sub>4</sub><sup>+</sup> generally increases plant growth.

Carbon dioxide and CH<sub>4</sub> emissions generally increased with increasing nutrients, but the effect was not statistically significant in either of our laboratory or field studies. We hypothesize there was not a significant nutrient effect on GHG emissions in the observational field study because the soils at each site differed significantly from each other, particularly in terms of soil OM content. We likewise did not see a nutrient effect in our controlled laboratory study, likely due to high initial concentrations of NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup>, and SRP in the soil porewater (1.28 mg/g, 94.2 mg/g, and 13.5 mg/g, respectively). Since the soil microbes had access to this nutrient-rich porewater under all treatment conditions, the N and P added within the surface water were less important as these nutrients were already likely to be non-limiting.

## 5. CONCLUSION


The emissions of CO<sub>2</sub> and CH<sub>4</sub> from treatment wetland soils increased with increasing temperatures between 25 and 35°C. The same general trend of increasing GHG production with increasing temperature was also observed in the field study, but the effect was not significant. Bioavailable nutrients in the surface water (NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, and SRP) did not show any statistical significance when considering GHG emissions

for any of the studies conducted; however, general increases in emissions were observed with increasing nutrients for each study. We suspect we did not observe a nutrient effect on GHG emissions for the field study because differences in soil properties (specifically, OM content) were so significant that surface water nutrient concentrations were not the limiting factor for soil microbial activity. Similarly, we did not see a nutrient effect in the controlled laboratory study due to the soils already being highly eutrophic, as exemplified by high porewater nutrient concentrations. In order to isolate the effect of bioavailable nutrients on GHG emissions from wetland soils, another laboratory study should be conducted with oligotrophic soils in order to isolate the effects of both nutrient concentration and temperature.


According to our findings, microbial communities that have  $\text{CO}_2$  as a byproduct of respiration may have a temperature threshold around 30–35°C, where metabolic processes are not stimulated at or above this range. This finding can be used to make accurate predictions for future climate change models (specifically for the C budget) when trying to understand the C flux from wetlands under climate change scenarios. Alternatively, a temperature threshold did not exist for  $\text{CH}_4$ , showing that this microbial community may have increased metabolic rates as temperature continues to rise. Though  $\text{CO}_2$  was still the predominant GHG being produced for both studies and all temperature and nutrient treatments, the global warming potential of  $\text{CH}_4$  must be taken into consideration when making predictions for future atmospheric temperatures. Another study should be conducted in order to identify different microbial communities within the soil, using qPCR before and after an extended incubation period at the same temperature and nutrient treatments selected in the controlled laboratory study to determine how these communities change under both treatments.




Figure 1: Schematic of the OWP indicating cell numbers and flow trains.




**Figure 2:** Carbon dioxide flux for the observational field study for both seasons and temperature treatments. Error bars represent standard error of the mean. Ambient and elevated temperatures for winter were 22°C and 26°C, respectively. Ambient and elevated temperatures for summer were 32°C and 36°C, respectively. Error bars with the same letter are not significantly different.




**Figure 3:** Methane flux for the observational field study for both seasons and temperature treatments. Ambient and elevated temperatures for winter were 22°C and 26°C, respectively. Ambient and elevated temperatures for summer were 32°C and 36°C, respectively. Error bars represent standard error of the mean. Error bars with the same letter are not significantly different.



**Figure 4a:** Carbon dioxide flux for the controlled laboratory study with respect to the added nutrient concentration. Error bars represent standard error of the mean. Error bars with the same letter are not significantly different.



**Figure 4b:** Methane flux for the controlled laboratory study with respect to the added nutrient concentration. Error bars represent standard error of the mean. Error bars with the same letter are not significantly different.



**Table 1:** Electron acceptors used for microbial respiration and their end products. Metabolic energy yield decreases with each subsequent pathway.

<i>Sequence of Reduction</i>	<i>Electron Acceptor</i>	<i>Electron Donor</i>	<i>End Products</i>
1	O <sub>2</sub>	Organic Matter	CO <sub>2</sub> + H <sub>2</sub> O
2	NO <sub>3</sub> <sup>-</sup>	Organic Matter	CO <sub>2</sub> + N <sub>2</sub> O/N <sub>2</sub>
3	Mn <sup>4+</sup>	Organic Matter	CO <sub>2</sub> + Mn <sup>2+</sup>
4	Fe <sup>3+</sup>	Organic Matter	CO <sub>2</sub> + Fe <sup>2+</sup>
5	SO <sub>4</sub> <sup>2-</sup>	Organic Matter	CO <sub>2</sub> + H <sub>2</sub> S, S
6	CO <sub>2</sub> /H <sub>2</sub>	Organic Matter	CH <sub>4</sub>

**Table 2:** Nutrient concentrations added to 1 L of nanopure water to mimic surface water nutrient concentrations.

	<i>High</i>	<i>Medium</i>	<i>Low</i>
<i>Phosphate (mL)</i>	0.4	0.2	0
<i>Ammonium (mL)</i>	1.1	0.55	0
<i>Glucose (g)</i>	0.146	0.145	0.144

**Table 3:** Correlations of soil physiochemical properties, soil/water nutrients, and MBC to both GHG emissions for both seasons of the observational field study. *Italic values denote significance. Critical value is 0.497 and n = 16.*

	CO <sub>2</sub> Flux	CH <sub>4</sub> Flux
CO <sub>2</sub> Flux		
CH <sub>4</sub> Flux	<i>0.712</i>	
TP (mg/g)	-0.292	-0.277
TN (%)	<i>0.812</i>	<i>0.915</i>
TC (%)	<i>0.800</i>	<i>0.912</i>
C/N Ratio	-0.595	-0.569
MBC	<i>0.873</i>	<i>0.937</i>
pH	0.207	0.249
Organic Matter	<i>0.707</i>	<i>0.878</i>
Bulk Density	-0.732	-0.834
Moisture Content	<i>0.758</i>	<i>0.850</i>
Surface Water NO <sub>3</sub> <sup>-</sup>	-0.351	-0.299
Surface Water NH <sub>4</sub>	0.221	<i>0.558</i>
Surface Water SRP	-0.225	0.197
Porewater NO <sub>3</sub> <sup>-</sup>	0.255	0.477
Porewater NH <sub>4</sub>	0.427	<i>0.669</i>
Porewater SRP	-0.449	-0.282

## REFERENCES

1. Andersen, J.M. 1976. An ignition method for determination of total phosphorus in lake sediments. *Water Res* 10:329–331. doi: 10.1016/0043-1354(76)90175-5.
2. Ansari, A.A. 2011. *Eutrophication: causes, consequences and control*. Dordrecht; New York: Springer.
3. Ballantyne, A. P., C.B Alden., J.B. Miller, P.P. Tans, and J.W. White. 2012. Increase in observed net carbon dioxide uptake by land and oceans during the last 50 years. *Nature*, 488, 70–72.
4. Bodelier, P.L., Laanbroek, H.J. 2004. Nitrogen as a regulatory factor of methane oxidation in soils and sediments. *FEMS Microbiol. Ecol.* 47, 265–277.
5. Bridgham SD, Richardson C.J. 1992. Mechanisms controlling soil respiration (CO<sub>2</sub> and CH<sub>4</sub>) in southern peatlands. *Soil Biol Biochem* 24:1089–1099
6. Brooks, P.D., Schmidt, S.K., Williams, M.W. 1997. Winter production of CO<sub>2</sub> and N<sub>2</sub>O from alpine tundra: environmental controls and relationship to inter-system C and N fluxes. *Oecologia* 110, 403–413.
7. Butterbach-Bahl, K., Kiese, R., Liu, C.Y. 2011. Measurement of biosphere-atmosphere exchange of CH<sub>4</sub> in terrestrial ecosystems. *Meth. Enzymol.* 495, 271–287.
8. Chapuis-Lardy, L., Wrage, N., Metay, A., Chotte, J.L., Bernoux, M. 2007. Soils, a sink for N<sub>2</sub>O? A review. *Glob. Change Biol.* 13, 1–17.
9. Chen, H. 2011. Surface-Flow Constructed Treatment Wetlands for Pollutant Removal: Applications and Perspectives. *Wetlands*, 31(4), 805-814.
10. Chin K.J., Lukow T., Conrad R. 1999. Effect of temperature on structure and function of the methanogenic archaeal community in an anoxic rice field soil. *Appl Environ Microbiol* 65:2341–2349.
11. Ciais, P., C. Sabine, G. Bala, L. Bopp, V. Brovkin, J. Canadell, A. Chhabra, R. DeFries, J. Galloway, M. Heimann, C. Jones, C. Le Quéré, R.B. Myneni, S. Piao and P. Thornton. 2013. Carbon and Other Biogeochemical Cycles. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
12. Dalal, R.C., Allen, D.E. 2008. Greenhouse gas fluxes from natural ecosystems. Turner review no. 18, *Australian J. Bot.* 56, 369–407.
13. Dunfield, P., Knowles, R. 1995. Kinetics of inhibition of methane oxidation by nitrate, nitrite, and ammonium in a humisol. *Appl. Environ. Microbiol.* 61, 3129–3135.
14. Fang C, Moncrieff J.B. 2001. The dependence of soil CO<sub>2</sub> efflux on temperature. *Soil Biol Biochem* 33:155–165.
15. Food and Agriculture Organization (FAO) of the United Nations. 2000. "Fertilizer requirements in 2015 and 2030." Rome: FAO.
16. Gulledge, J., Schimel, J.P. 1998. Low-concentration kinetics of atmospheric CH<sub>4</sub> oxidation in soil and mechanism of NH<sub>4</sub><sup>+</sup> inhibition. *Appl. Environ. Microbiol.* 64, 4291–4298.
17. Hagner, M., Kemppainen, R., Jauhiainen, L., Tiilikkala, K., & Setälä, H. 2016. The effects of birch (*Betula* spp.) biochar and pyrolysis temperature on soil properties and plant growth. *Soil & Tillage Research*, 224. doi:10.1016/j.still.2016.06.006
18. Holst, J., Liu, C., Yao, Z., Brüggemann, N., Zheng, X., Giese, M., Butterbach-Bahl, K., 2008. Fluxes of nitrous oxide, methane and carbon dioxide during freezing–thawing cycles in an Inner Mongolian steppe. *Plant Soil* 308, 105–117.
19. Houghton, J.T. 2001. "Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press. p. 881.
20. Inglett, K.S., P.W. Inglett, K.R. Reddy, and T.Z. Osborne. 2012. Temperature sensitivity of greenhouse gas production in wetland soils of different vegetation. *Biogeochemistry*.

21. IPCC, 2007: Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, M.L. Parry, O.F. Canziani, J.P. Palutikof, P.J. van der Linden and C.E. Hanson, Eds., Cambridge University Press, Cambridge, UK, 976pp
22. IPCC, 2007: Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, Pachauri, R.K and Reisinger, A. (eds.)]. IPCC, Geneva, Switzerland, 104 pp.
23. Johnson, J.M.F., A.J. Franzluebbers, S.L. Weyers, and D.C. Reicosky, 2007. Agricultural opportunities to mitigate greenhouse gas emissions. *Environ. Pollut.* 150(1): 107–124.
24. Kadlec, R. H., Wallace, S. 2008. Treatment wetlands—Second Edition. CRC press.
25. Kadlec, R.H., 2009. Comparison of free water and horizontal subsurface treatment wetlands. *Ecol. Eng.* 35, 159-174.
26. Larmola, T., Bubier, J.L., Kobyljanec, C., Basiliko, N., Juutinen, S., Humphreys, E., and Moore, T. R. 2013. Vegetation feedbacks of nutrient addition lead to a weaker carbon sink in an ombrotrophic bog. *Global Change Biology*, (12), 3729.
27. Ludwig, J., Meixner, F.X., Vogel, B., Förstner, J. 2001. Soil-air exchange of nitric oxide: an overview of processes, environmental factors, and modeling studies. *Biogeochemistry* 52, 225–257.
28. Marimon, Z. A., & Chang, N. 2016. Evaluating floating treatment wetlands to improve nitrogen removal in a wet detention pond. Orlando, Florida: University of Central Florida, 2016.
29. Martinez, C.J, Wise, W.R. 2003. Hydraulic Analysis of Orlando Easterly Wetland. *Journal Of Environmental Engineering*, 129(6), 553.
30. McCormick, P. V., and J.A. Laing. 2003. Effects of increased phosphorus loading on dissolved oxygen in a subtropical wetland, the Florida Everglades. *Wetl. Ecol. Manag.* 11(3): 199–215.
31. McKenzie C, Schiff S, Aravena R, Kelly C, St. Louis V, 1998. Effect of temperature on production of CH<sub>4</sub> and CO<sub>2</sub> from Peat in a Natural and Flooded Boreal Forest Wetland. *Clim Change* 40:247–266
32. National Oceanic and Atmospheric Administration (NOAA), 2017. “Global Climate Change: Carbon Dioxide.” Retrieved from <https://climate.nasa.gov/vital-signs/carbon-dioxide/>
33. Niu, S., Wu, M., Han, Y.I., Xia, J., Zhang, Z.H., Yang, H., Wan, S. 2010. Nitrogen effects on net ecosystem carbon exchange in a temperate steppe. *Glob. Change Biol.* 16, 144–155.
34. O'Connel, A.M. 1990. Microbial decomposition (respiration) of litter in eucalypt forests of south-western Australia: an empirical model based on laboratory incubations. *Soil Biology and Biochemistry* 22, 153–160.
35. Oertel, C.J. Matschullat, K. Zurba, F. Zimmermann, and S. Erasmi. 2016. Greenhouse gas emissions from soils: A review. *Chemie der Erde–Geochemistry*.
36. IPCC, 2007: Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, M.L. Parry, O.F. Canziani, J.P. Palutikof, P.J. van der Linden and C.E. Hanson, Eds., Cambridge University Press, Cambridge, UK, 976pp.
37. Peng, Q., Dong, Y., Qi, Y., Xiao, S., He, Y., Ma, T. 2011. Effects of nitrogen fertilization on soil respiration in temperate grassland in Inner Mongolia, China. *Environ. Earth Sci.* 62, 1163–1171.
38. Reddy KR, DeLaune RD. 2008. Biogeochemistry of Wetlands: Science and Applications. CRC Press
39. Rothfeld, K. 2015. Orlando Easterly Wetlands 2015 Annual Report. Retrieved from <http://www.cityoforlando.net/wetlands/wp-content/uploads/sites/59/2016/07/Annual-Report-2015.pdf>

40. Schimel, J. 2000. Rice, microbes and methane. *Nature* 403, 375–377.

41. Sees M, Turner D. 1997. “Orlando easterly wetlands; Management of a constructed wetland treatment system.” Proc., Florida Lake Management Society 1997 Annual Meeting, May 7–9, 1997 Palm Beach Shores, Fla. <http://www.ces.fau.edu/library/flms/24.html>.

42. Shi, W.Y., Yan, M.J., Zhang, J.G., Guan, J.H., Du, S. 2014. Soil CO<sub>2</sub> emissions from five different types of land use on the semiarid Loess Plateau of China, with emphasis on the contribution of winter soil respiration. *Atmos. Environ.* 88, 74–82.

43. Sonstebj, A., Solhaug, K.A., and Heide, O.M. 2016. Functional growth analysis of 'Sonata' strawberry plants grown under controlled temperature and daylength conditions. *Scientia Horticulturae*, 26. doi:10.1016/j.scienta.2016.08.003

44. U.S. Environmental Protection Agency (U.S. EPA). 1993. Methods for the determination of inorganic substances in environmental samples. EPA/600/R-93/100. Washington, DC.

45. U.S. Environmental Protection Agency (U.S. EPA). 1993. “Wetland treatment systems: A Case history, The Orlando easterly wetlands reclamation project.” *EPA832-R-93-005i*, U.S. EPA Office of Water, Washington, D.C.

46. United Nations Population Division (UNPD). 2008. World Population Prospectus: the 2008 revision population database. Available at: <http://esa.un.org/unpp/>.

47. Vance E.D., Brookes P.C., Jenkinson D.S. 1987 An extraction method for measuring soil microbial biomass C. *Soil Biol Biochem* 19:703–707. doi: 10.1016/0038-0717(87)90052-6

48. Webber, C.D. 2010. Eutrophication: Ecological Effects, Sources, Prevention and Reversal. New York: Nova Science Pub. Inc.,; 2010.